

**MUGARD (carbomer; glycerin; polysorbate)**

Effective Date: 1/28/14

Date Developed: 1/28/14 by Robert Sterling, MD

Last Approval Date: 1/26/16, 1/24/17, 1/23/18, 1/22/19, 2/18/20

Mugard is an oral rinse which acts as a mucoadhesive and forms a protective coating over the oral mucosa

**Authorization:** For the management of oral mucositis/stomatitis and all types of oral wounds, including aphthous ulcers/canker sores and traumatic ulcers.

**Dosing:** Pour 5mL into mouth and rinse entire oral cavity for 30–60 seconds, or longer. May use up to 10mL to fully coat inside of mouth. After rinsing, expel or swallow. Repeat 4–6 times/day. Avoid eating or drinking at least 1 hour after rinsing. Not recommended for children.

**PRECAUTIONS:** Do not dilute. Avoid food or liquids for at least one hour after use

**DRUG INTERACTIONS:** none known

**REFERENCES**

Sonis et al, Cancer, 2000, 89:2103

Epstein et al, Cancer, 2000, 89:2258

**Revision History:**

Date Reviewed/No Updates: 1/13/15 by C. Sanders, MD

Date Approved by P&T Committee: 1/27/15

Date Reviewed/No Updates: 1/26/16 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/26/16

Date Reviewed/No Updates: 1/24/17 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/24/17

Date Reviewed/No Updates: 1/23/18 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/23/18

Date Reviewed/No Updates: 1/22/19 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/22/19

Date Reviewed/No Updates: 2/18/20 by H. Taekman, MD; R. Sterling, MD

Date Approved by P&T Committee: 2/18/20

<b>Revision Date</b>	<b>Content Revised (Yes/No)</b>	<b>Contributors</b>	<b>Review/Revision Notes</b>
1/24/17	No	Catherine Sanders, MD; Robert Sterling, MD	Annual review
1/23/18	No	Catherine Sanders, MD; Robert Sterling, MD	Annual review
1/22/19	No	Catherine Sanders, MD; Robert Sterling, MD	Annual review
2/18/20	No	Howard Taekman, MD; Robert Sterling, MD	Annual review