

RILUTEK (riluzole)

Effective Date: 1/28/14

Date Developed: 1/28/14 by Robert Sterling, MD

Last Approval Date: 1/26/16, 1/24/17, 1/23/18, 1/22/19, 2/18/20

Rilutek is used in the palliative treatment of amyotrophic lateral sclerosis (ALS) to extend survival and/or time to tracheostomy

Authorization: patients diagnosed with ALS

Dosing: 50mg every 12 hours, one hour before or two hours after a meal; little or no benefit to increased doses

PRECAUTIONS: use with caution in patients with liver disease; neutropenia; hypersensitivity pneumonitis; nausea; abdominal pain; constipation or diarrhea; methemoglobinemia; asthenia

DRUG INTERACTIONS: none reported

REFERENCES

Bensimon G, Lacomblez L, Meininger V, et al, "A Controlled Trial of Riluzole in Amyotrophic Lateral Sclerosis. ALS/Riluzole Study Group," N Engl J Med, 1994, 330(9):585-91

Wokke J, "Riluzole," Lancet, 1996, 348(9030):795-9.

Revision History:

Date Reviewed/No Updates: 1/13/15 by C. Sanders, MD

Date Approved by P&T Committee: 1/27/15

Date Reviewed/No Updates: 1/26/16 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/26/16

Date Reviewed/No Updates: 1/24/17 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/24/17

Date Reviewed/No Updates: 1/23/18 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/23/18

Date Reviewed/No Updates: 1/22/19 by C. Sanders, MD; R. Sterling, MD

Date Approved by P&T Committee: 1/22/19

Date Reviewed/No Updates: 2/18/20 by H. Taekman, MD; R. Sterling, MD

Date Approved by P&T Committee: 2/18/20

Revision Date	Content Revised (Yes/No)	Contributors	Review/Revision Notes
1/24/17	No	Catherine Sanders, MD; Robert Sterling, MD	Annual review
1/23/18	No	Catherine Sanders, MD; Robert Sterling, MD	Annual review
1/22/19	No	Catherine Sanders, MD; Robert Sterling, MD	Annual review
2/18/20	No	Howard Taekman, MD; Robert Sterling, MD	Annual review