

POLICY: Inflammatory Conditions – Ilumya™ (tildrakizumab-asmn for subcutaneous injection – Sun Pharmaceuticals)

DATE REVIEWED: 04/22/2020

OVERVIEW

Ilumya is a humanized immunoglobulin G monoclonal antibody that binds to interleukin (IL)-23, a pro-inflammatory cytokine.¹ It binds to the p19 subunit of IL-23 and inhibits the intracellular and downstream signaling of IL-23 which is required for the terminal differentiation and survival of T helper 17 cells. Ilumya is indicated for treatment of adults with moderate to severe plaque psoriasis who are candidates for systemic therapy or phototherapy. It is administered subcutaneously (SC) at Weeks 0 and 4 and then once every 12 weeks thereafter. Ilumya should be administered by a healthcare professional. Safety and efficacy have not been established in patients < 18 years of age.

Disease Overview

Although the etiology of psoriasis is not fully established, abnormal keratin formation, epidermal proliferation, activation of the immune system, and hereditary factors appear to play roles in the pathogenesis of the disease. In psoriasis, levels of IL-23p40 and IL-12/23p40 messenger RNA are upregulated but decrease with treatment. By blocking the release of proinflammatory cytokines and chemokines, Ilumya has an inhibitory effect on the inflammatory process.

Guidelines

Joint guidelines from the American Academy of Dermatology and National Psoriasis Medical Board (2019) have been published for management of psoriasis with biologics.² These guidelines list Ilumya as a monotherapy treatment option for patients with moderate to severe plaque psoriasis. Guidelines from the European Dermatology Forum (2015) recommend biologics (i.e., etanercept, adalimumab, infliximab, Stelara SC) as second-line therapy for induction and long-term treatment if phototherapy and conventional systemic agents have failed, are contraindicated, or are not tolerated.³

POLICY STATEMENT

Prior authorization is recommended for medical benefit coverage of Ilumya. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication(s). Extended approvals are allowed if the patient continues to meet the criteria and dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

Because of the of the specialized skills required for evaluation and diagnosis of patients treated with Ilumya, as well as the monitoring required for adverse events and long-term efficacy, initial approval requires Ilumya to be prescribed by or in consultation with a physician who specializes in the condition being treated.

RECOMMENDED AUTHORIZATION CRITERIA

FDA-Approved Indication

- i. Plaque Psoriasis.** Approve for the duration noted if the patient meets ONE of the following (A or B):
- A) Initial Therapy.** Approve for 3 months if the patient meets ALL of the following criteria (i, ii, and iii):
- i.** The patient is ≥ 18 years of age; AND
 - ii.** The patient meets ONE of the following conditions (a or b):
 - a)** The patient has tried at least at least one traditional systemic agent for psoriasis for at least 3 months, unless intolerant.
Note: Examples of one traditional systemic agent include methotrexate (MTX), cyclosporine, acitretin tablets, or psoralen plus ultraviolet A light (PUVA). An exception to the requirement for a trial of one traditional systemic agent for psoriasis can be made if the patient has already had a 3-month trial or previous intolerance to at least one biologic Refer to [Appendix](#) for examples of biologics used for plaque psoriasis. These patients who have already tried a biologic for psoriasis are not required to “step back” and try a traditional systemic agent for psoriasis); OR
 - b)** The patient has a contraindication to methotrexate (MTX), as determined by the prescriber; AND
 - iii.** Ilumya is prescribed by or in consultation with a dermatologist.
- B) Patient is Currently Receiving Ilumya.** Approve for 1 year if the patient has responded, as determined by the prescriber.
Note: The patient may not have a full response, but there should have been a recent or past response to Ilumya.

Dosing. Approve the following dosing (A and B):

- A)** The dose is 100 mg/kg given as a subcutaneous injection; AND
- B)** Doses are administered at Weeks 0 and 4, then not more frequently than once every 12 weeks thereafter.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Ilumya has not been shown to be effective, or there are limited or preliminary data or potential safety concerns that are not supportive of general approval for the following conditions. Rationale for non-coverage for these specific conditions is provided below. (Note: This is not an exhaustive list of Conditions Not Recommended for Approval.)

- 1. Concurrent Use with other Biologics or with Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs).** Data are lacking evaluating concomitant use of Ilumya with another biologic or with a targeted synthetic DMARD for an inflammatory condition (see [Appendix](#) for examples). Combination therapy with biologics and/or biologics + targeted synthetic DMARDs has a potential for a higher rate of adverse effects and lack controlled trial data in support of additive efficacy.⁴

Note: This does NOT exclude the use of MTX (a traditional systemic agent used to treat psoriasis) in combination with Ilumya.

2. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Ilumya™ injection [prescribing information]. Whitehouse Station, NJ: Sun Pharmaceuticals; October 2019.
2. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
3. Nast A, Gisondi P, Ormerod AD, et al. European S3-Guidelines on the systemic treatment of psoriasis vulgaris – Update 2015 – Short version – EDF in cooperation with EADV and IPC. *J Eur Acad Dermatol Venereol.* 2015;29(12):2277-2294.
4. Reich K, Papp KA, Blauvelt A, et al. Tildrakizumab versus placebo or etanercept for chronic plaque psoriasis (reSURFACE 1 and reSURFACE 2): results from two randomised controlled, phase 3 trials. *Lancet.* 2017;390(10091):276-288.

HISTORY

Type of Revision	Summary of Changes	Date Reviewed
New Policy	--	04/22/2020

APPENDIX

	Mechanism of Action	Examples of Inflammatory Indications for Products*
Biologics		
Adalimumab SC Products (Humira®, biosimilars)	Inhibition of TNF	AS, CD, PJIA, PsO, PsA, RA, SJIA, UC
Cimzia® (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, PsO, PsA, RA
Etanercept SC Products (Enbrel®, biosimilars)	Inhibition of TNF	AS, PJIA, PsO, PsA, RA, SJIA
Infliximab IV Products (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PJIA, PsO, PsA, RA, SJIA, UC
Simponi®, Simponi® Aria™ (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC IV formulation: AS, PsA, RA
Actemra® (tocilizumab IV infusion, tocilizumab SC injection)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA IV formulation: PJIA, RA, SJIA
Kezara® (sarilumab SC injection)	Inhibition of IL-6	RA
Orencia® (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: PJIA, PSA, RA IV formulation: PJIA, PsA, RA
Rituximab IV Products (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
Ilaris (canakinumab SC injection)	Inhibition of IL-1β	SJIA
Kineret® (anakinra SC injection)	Inhibition of IL-1	RA, SJIA [^]
Stelara® (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC IV formulation: CD, UC
Siliq™ (brodalumab SC injection)	Inhibition of IL-17	PsO
Cosentyx™ (secukinumab SC injection)	Inhibition of IL-17A	AS, PsO, PsA
Taltz® (ixekizumab SC injection)	Inhibition of IL-17A	AS, PsO, PsA
Ilumya™ (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
Skyrizi™ (risankizumab-rzza SC injection)	Inhibition of IL-23	PsO
Tremfya™ (guselkumab SC injection)	Inhibition of IL-23	PsO
Entyvio™ (vedolizumab IV infusion)	Integrin receptor antagonist	CD, UC
Targeted Synthetic DMARDs		
Otezla® (apremilast tablets)	Inhibition of PDE4	PsO, PsA
Olumiant® (baricitinib tablets)	Inhibition of the JAK pathways	RA
Rinvoq® (upadacitinib extended-release tablets)	Inhibition of the JAK pathways	RA
Xeljanz®, Xeljanz XR (tofacitinib tablets, tofacitinib extended-release tablets)	Inhibition of the JAK pathways	RA, PsA, UC

* Not an all-inclusive list of indication (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; IV – Intravenous, IL – Interleukin; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AS – Ankylosing spondylitis; CD – Crohn’s disease; PJIA – Polyarticular juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; SJIA – Sytemice juvenile idiopathic arthritis; UC – Ulcerative colitis. [^] Off-label use of SJIA supported in guidelines.